arXiv:1111.4173v1 [math.DG] 17 Nov 2011

The local description of the Ricci and Bianchi identities for an h-normal N-linear connection on the dual 1-jet space $J^{1*}(\mathcal{T}, M)$

Alexandru Oană and Mircea Neagu

Abstract

In this paper we describe the local Ricci and Bianchi identities for an h-normal N-linear connection $D\Gamma(N)$ on the dual 1-jet space $J^{1*}(\mathcal{T}, M)$. To reach this aim, we firstly give the expressions of the local distinguished (d-) adapted components of torsion and curvature tensors produced by $D\Gamma(N)$, and then we analyze their attached local Ricci identities. The derived deflection d-tensor identities are also presented. Finally, we expose the local expressions of the Bianchi identities (in the particular case of an h-normal N-linear connection of Cartan type), which geometrically connect the local torsion and curvature d-tensors of the linear connection $D\Gamma(N)$.

2000 Mathematics Subject Classification: 53B40, 53C60, 53C07.

Key words and phrases: dual 1-jet spaces, nonlinear connections, h-normal N-linear connections of Cartan type, Ricci and Bianchi identities.

1 Introduction

According to Olver's opinion [21], we consider that the 1-jet spaces and their duals are natural houses for the study of classical and quantum field theories. For such a reason, the differential geometry of 1-jet spaces was intensively studied, in a contravariant approach, by a lot of authors: Saunders [22], Asanov [1], Neagu and Udrişte (see [18], [19], [20]), and many others.

In the last decades, numerous physicists and geometers were preoccupied by the development of that so-called the *covariant Hamiltonian geometry of physical fields*, which is a multi-parameter, or multi-time, extension of the classical Hamiltonian formulation from Mechanics. In such a perspective, we point out that the covariant Hamiltonian geometry of physical fields appears in the literature of specialty in three distinct variants: **(1)** ► the *multisymplectic geometry* − developed by Gotay, Isenberg, Marsden, Montgomery and their co-workers (see [9], [8]) on a finite-dimensional multisymplectic phase space; **(2)** ► the *polysymplectic geometry* − elaborated by Giachetta, Mangiarotti and Sardanashvily (see [6], [7]), which emphasizes the relations between the equations of first order Lagrangian field theory on fiber bundles and the covariant Hamilton equations on a finite-dimensional polysymplectic phase space; **(3)** ▶ the *De Donder-Weyl Hamiltonian geometry* – studied by Kanatchikov (see [10], [11], [12]) as opposed to the conventional field-theoretical Hamiltonian formalism, which requires the space + time decomposition and leads to the picture of a field as a mechanical system with infinitely degrees of freedom.

From a geometrical point of view, following the ideas initially stated by Asanov [1], a multi-time Lagrange contravariant geometry on 1-jet spaces (in the sense of d-linear connections, d-torsions and d-curvatures) was recently developed by Neagu and Udrişte in [18], [19] and [20]. This 1-jet geometrical theory is a natural multi-time extension of the classical Lagrangian geometry on tangent bundles, initiated and developed by Miron and Anastasiei [14].

On the other hand, suggested by the field theoretical extension of the basic structures of classical Analytical Mechanics within the framework of the De Donder-Weyl covariant Hamiltonian formulation, the geometrical studies of Miron [13], Atanasiu [3], [2] and others led to the development of the Hamilton geometry on cotangent bundles, which is synthesized in the book [15]. Note that the Miron-Atanasiu Hamiltonian geometrical ideas from cotangent bundles represent the point start for the development of the jet covariant Riemann-Hamilton geometry depending on polymomenta, which is presented in the Atanasiu-Neagu papers [4] and [5]. In this paper we are going on the jet multi-time Hamiltonian geometrical studies from [4] and [5].

2 Components of N-linear connections on dual 1-jet bundle $J^{1*}(\mathcal{T}, M)$

Let $\mathcal{T}$ and M be a *temporal* (resp. *spatial*) real, smooth manifold of dimension m (resp. n), whose coordinates are $(t^a)_{a=\overline{1,m}}$, respectively $(x^i)_{i=\overline{1,n}}$. Note that, throughout this paper, the indices a, b, c, ... run from 1 to m, while the indices i, j, k, ... run from 1 to n. The Einstein convention of summation is also adopted all over this work.

Let $J^{1*}(\mathcal{T}, M)$ be the dual 1-jet fibre bundle, whose coordinates (t^a, x^i, p^a_i) are induced from $\mathcal{T}$ and M. The coordinate transformations from the product manifold $\mathcal{T}\times\ M$ produce on $J^{1*}(\mathcal{T}, M)$ the following coordinate transformations:

$$\widetilde{t}^a = \widetilde{t}^a\left(t^b\right), \quad \widetilde{x}^i = \widetilde{x}^i\left(x^j\right), \quad \widetilde{p}^a_i = \frac{\partial x^j}{\partial \widetilde{x}^i}\frac{\partial \widetilde{t}^a}{\partial t^b}p^b_j,$$

where $\det\left(\partial\widetilde{t}^a/\partial t^b\right) \neq 0$ and $\det\left(\partial\widetilde{x}^i/\partial x^j\right) \neq 0$.

Definition 2.1 *A pair of local functions on $E^* = J^{1*}(\mathcal{T}, M)$, denoted by*

$$N = \left(\underset{1}{N}{}^{(a)}_{(i)b},\ \underset{2}{N}{}^{(a)}_{(i)j}\right),$$

whose local components obey the transformation rules

$$\underset{1}{\widetilde{N}}{}^{(b)}_{(j)c}\frac{\delta\widetilde{t}^c}{\delta t^a} = \underset{1}{N}{}^{(c)}_{(k)a}\frac{\delta\widetilde{t}^b}{\delta t^c}\frac{\partial x^k}{\partial\widetilde{x}^j} - \frac{\partial\widetilde{p}^b_j}{\partial t^a},$$

$$\underset{2}{\widetilde{N}}{}^{(b)}_{(j)k}\frac{\partial\widetilde{x}^k}{\partial x^i}=\underset{2}{N}{}^{(c)}_{(k)i}\frac{\delta\widetilde{t}^b}{\delta t^c}\frac{\partial x^k}{\partial\widetilde{x}^j}-\frac{\partial\widetilde{p}^b_j}{\partial x^i},$$

is called a ***nonlinear connection*** *on* E^*. *The components* $\underset{1}{N}{}^{(a)}_{(i)b}$ *(resp.* $\underset{2}{N}{}^{(a)}_{(i)j}$*) are called the* ***temporal*** *(resp.* ***spatial****)* ***components*** *of* N.

Example 2.2 *Let* $h_{ab}\left(t^f\right)$ *(resp.* $\varphi_{ij}\left(x^k\right)$*) be a semi-Riemannian metric on the temporal manifold* $\mathcal{T}$ *(resp. spatial manifold* M*). Taking into account the local transformation rules of the Christoffel symbols* $\chi^a_{bc}(t)$ *(resp.* $\Gamma^k_{ij}(x)$*) of the metrics* $h_{ab}(t)$ *(resp.* $\varphi_{ij}(x)$*), then the pair of local functions*

$$N_0=\left(\underset{1}{\overset{0}{N}}{}^{(a)}_{(i)b},\ \underset{2}{\overset{0}{N}}{}^{(a)}_{(i)j}\right),$$

where

$$\underset{1}{\overset{0}{N}}{}^{(a)}_{(i)b}=\chi^a_{bc}p^c_i,\qquad \underset{2}{\overset{0}{N}}{}^{(a)}_{(i)j}=-\Gamma^k_{ij}p^a_k,$$

represents a nonlinear connection on E^*. *This is called the* ***canonical nonlinear connection attached to the metrics*** $h_{ab}(t)$ ***and*** $\varphi_{ij}(x)$.

In what follows, we fix a nonlinear connection on E^*, and we consider the *adapted bases* of the nonlinear connection N, defined by

$$\left\{\frac{\delta}{\delta t^a},\frac{\delta}{\delta x^i},\frac{\partial}{\partial p^a_i}\right\}\subset\mathcal{X}\left(E^*\right),\quad \left\{dt^a,dx^i,\delta p^a_i\right\}\subset\mathcal{X}^*\left(E^*\right), \tag{2.1}$$

where

$$\begin{aligned}\frac{\delta}{\delta t^a}&=\frac{\partial}{\partial t^a}-\underset{1}{N}{}^{(b)}_{(j)a}\frac{\partial}{\partial p^b_j},\\ \frac{\delta}{\delta x^i}&=\frac{\partial}{\partial x^i}-\underset{2}{N}{}^{(b)}_{(j)i}\frac{\partial}{\partial p^b_j},\\ \delta p^a_i&=dp^a_i+\underset{1}{N}{}^{(a)}_{(i)b}dt^b+\underset{2}{N}{}^{(a)}_{(i)j}dx^j.\end{aligned}$$

It is important to note that the transformation rules of the elements of the adapted bases (2.1) are tensorial ones:

$$\begin{array}{l}\dfrac{\delta}{\delta t^a}=\dfrac{\partial\widetilde{t}^b}{\partial t^a}\dfrac{\delta}{\delta\widetilde{t}^b},\quad \dfrac{\delta}{\delta x^i}=\dfrac{\partial\widetilde{x}^j}{\partial x^i}\dfrac{\delta}{\delta\widetilde{x}^j},\quad \dfrac{\partial}{\partial p^a_i}=\dfrac{\partial\widetilde{t}^b}{\partial t^a}\dfrac{\partial x^i}{\partial\widetilde{x}^j}\dfrac{\partial}{\partial\widetilde{p}^b_j},\\[2ex] dt^a=\dfrac{\partial t^a}{\partial\widetilde{t}^b}d\widetilde{t}^b,\quad dx^i=\dfrac{\partial x^i}{\partial\widetilde{x}^j}d\widetilde{x}^j,\quad \delta p^a_i=\dfrac{\partial t^a}{\partial\widetilde{t}^b}\dfrac{\partial\widetilde{x}^j}{\partial x^i}\delta\widetilde{p}^b_j.\end{array} \tag{2.2}$$

Remark 2.3 *The simple tensorial transformation rules (2.2) of the adapted bases (2.1) determined us to describe in what follows all geometrical objects on the dual* 1*-jet space* $J^{1*}(\mathcal{T},M)$ *in adapted local components.*

In order to develop the geometrical theory of N-linear connections on the dual 1-jet space E^*, we need the following result:

Proposition 2.4 (i) *The Lie algebra* $\mathcal{X}(E^*)$ *of vector fields decomposes as*

$$\mathcal{X}(E^*) = \mathcal{X}(\mathcal{H}_{\mathcal{T}}) \oplus \mathcal{X}(\mathcal{H}_M) \oplus \mathcal{X}(\mathcal{V}),$$

where

$$\mathcal{X}(\mathcal{H}_{\mathcal{T}}) = \text{Span}\left\{\frac{\delta}{\delta t^a}\right\}, \quad \mathcal{X}(\mathcal{H}_M) = \text{Span}\left\{\frac{\delta}{\delta x^i}\right\}, \quad \mathcal{X}(\mathcal{V}) = \text{Span}\left\{\frac{\partial}{\partial p_i^a}\right\}.$$

(ii) *The Lie algebra* $\mathcal{X}^*(E^*)$ *of covector fields decomposes as*

$$\mathcal{X}^*(E^*) = \mathcal{X}^*(\mathcal{H}_{\mathcal{T}}) \oplus \mathcal{X}^*(\mathcal{H}_M) \oplus \mathcal{X}^*(\mathcal{V}),$$

where

$$\mathcal{X}^*(\mathcal{H}_{\mathcal{T}}) = \text{Span}\{dt^a\}, \quad \mathcal{X}^*(\mathcal{H}_M) = \text{Span}\{dx^i\}, \quad \mathcal{X}^*(\mathcal{V}) = \text{Span}\{\delta p_i^a\}.$$

Let us consider that $h_{\mathcal{T}}$, h_M (horizontal) and v (vertical) are the canonical projections of the above decompositions. In this context, we introduce the following geometrical concept:

Definition 2.5 *A linear connection* $D : \mathcal{X}(E^*) \times \mathcal{X}(E^*) \to \mathcal{X}(E^*)$ *is called an* N***-linear connection*** *on* E^* *if and only if* $Dh_{\mathcal{T}} = 0, Dh_M = 0$ *and* $Dv = 0$.

It is obvious that the local description of the N-linear connection D on E^* is accomplished by *nine* unique adapted components

$$\begin{aligned} D\Gamma(N) = \Big(& A_{bc}^a,\ A_{jc}^i,\ -A_{(i)(b)c}^{(a)(j)},\ H_{bk}^a,\ H_{jk}^i,\ -H_{(i)(b)k}^{(a)(j)}, \\ & C_{b(c)}^{a(k)},\ C_{j(c)}^{i(k)},\ -C_{(i)(b)(c)}^{(a)(j)(k)} \Big), \end{aligned} \tag{2.3}$$

which are locally defined by the relations:

$$\begin{aligned} D_{\frac{\delta}{\delta t^c}} \frac{\delta}{\delta t^b} &= A_{bc}^a \frac{\delta}{\delta t^a},\ D_{\frac{\delta}{\delta t^c}} \frac{\delta}{\delta x^j} = A_{jc}^i \frac{\delta}{\delta x^i},\ D_{\frac{\delta}{\delta t^c}} \frac{\partial}{\partial p_j^b} = -A_{(i)(b)c}^{(a)(j)} \frac{\partial}{\partial p_i^a}, \\ D_{\frac{\delta}{\delta x^k}} \frac{\delta}{\delta t^b} &= H_{bk}^a \frac{\delta}{\delta t^a},\ D_{\frac{\delta}{\delta x^k}} \frac{\delta}{\delta x^j} = H_{jk}^i \frac{\delta}{\delta x^i},\ D_{\frac{\delta}{\delta x^k}} \frac{\partial}{\partial p_j^b} = -H_{(i)(b)k}^{(a)(j)} \frac{\partial}{\partial p_i^a}, \\ D_{\frac{\partial}{\partial p_k^c}} \frac{\delta}{\delta t^b} &= C_{b(c)}^{a(k)} \frac{\delta}{\delta t^a},\ D_{\frac{\partial}{\partial p_k^c}} \frac{\delta}{\delta x^j} = C_{j(c)}^{i(k)} \frac{\delta}{\delta x^i},\ D_{\frac{\partial}{\partial p_k^c}} \frac{\partial}{\partial p_j^b} = -C_{(i)(b)(c)}^{(a)(j)(k)} \frac{\partial}{\partial p_i^a}. \end{aligned}$$

Example 2.6 *Let* $N_0 = \left(\overset{0}{\underset{1}{N}}{}_{(i)b}^{(a)},\ \overset{0}{\underset{2}{N}}{}_{(i)j}^{(a)}\right)$ *be the canonical nonlinear connection produced by the semi-Riemannian metrics* (h_{ab}, φ_{ij}). *Taking into account the transformation rules of the Christoffel symbols* χ_{bc}^a *and* Γ_{jk}^i, *by local computations, we can show that the local components*

$$B\Gamma(N_0) = \left(\chi_{bc}^a,\ 0,\ -A_{(i)(b)c}^{(a)(j)},\ 0,\ \Gamma_{jk}^i,\ -H_{(i)(b)k}^{(a)(j)},\ 0,\ 0,\ 0\right)$$

where

$$A^{(a)(j)}_{(i)(b)c} = -\delta^j_i \chi^a_{bc}, \quad H^{(a)(j)}_{(i)(b)k} = \delta^a_b \Gamma^j_{ik},$$

verify the transformation rules of the components of an N-linear connection (for more details, see [5]*). Consequently, $B\Gamma\left(N_0\right)$ is an N_0-linear connection on E^*, which is called the* ***Berwald connection*** *of the metric pair $\left(h_{ab}, \varphi_{ij}\right)$.*

Now, let $D\Gamma(N)$ be an N-linear connection on E^*, locally defined by (2.3). The linear connection $D\Gamma(N)$ induces a linear connection on the set of d-tensors on the dual 1-jet fibre bundle $E^* = J^{1*}\left(\mathcal{T}, M\right)$, in a natural way. Thus, starting with a d-vector field X and a d-tensor field T, locally expressed by

$$\begin{aligned} X &= X^a \frac{\delta}{\delta t^a} + X^i \frac{\delta}{\delta x^i} + X^{(a)}_{(i)} \frac{\partial}{\partial p^a_i}, \\ T &= T^{ai(k)(d)\ldots}_{cj(b)(l)\ldots} \frac{\delta}{\delta t^a} \otimes \frac{\delta}{\delta x^i} \otimes \frac{\partial}{\partial p^d_l} \otimes dt^c \otimes dx^j \otimes \delta p^b_k \otimes \ldots, \end{aligned}$$

we can define the covariant derivative

$$\begin{aligned} & D_X T = X^g D_{\frac{\delta}{\delta t^g}} T + X^s D_{\frac{\delta}{\delta x^s}} T + X^{(g)}_{(s)} D_{\frac{\partial}{\partial p^g_s}} T = \\ & = \left\{ X^g T^{ai(k)(d)\ldots}_{cj(b)(l)\ldots/g} + X^s T^{ai(k)(d)\ldots}_{cj(b)(l)\ldots|s} + \right. \\ & \left. + X^{(g)}_{(s)} T^{ai(k)(d)\ldots}_{cj(b)(l)\ldots}\,|^{(s)}_{(g)} \right\} \frac{\delta}{\delta t^a} \otimes \frac{\delta}{\delta x^i} \otimes \frac{\partial}{\partial p^d_l} \otimes dt^c \otimes dx^j \otimes \delta p^b_k \otimes \ldots, \end{aligned}$$

where

- the $\mathcal{T}$*-horizontal covariant derivative* of $D\Gamma(N)$:

$$(h_{\mathcal{T}}) \quad \begin{cases} T^{ai(k)(d)\ldots}_{cj(b)(l)\ldots/g} = \dfrac{\delta T^{ai(k)(d)\ldots}_{cj(b)(l)\ldots}}{\delta t^g} + T^{fi(k)(d)\ldots}_{cj(b)(l)\ldots} A^a_{fg} + \\ + T^{ar(k)(d)\ldots}_{cj(b)(l)\ldots} A^i_{rg} + T^{ai(r)(d)\ldots}_{cj(f)(l)\ldots} A^{(f)(k)}_{(r)(b)g} + \ldots - \\ - T^{ai(k)(d)\ldots}_{fj(b)(l)\ldots} A^f_{cg} - T^{ai(k)(d)\ldots}_{cr(b)(l)\ldots} A^r_{jg} - T^{ai(k)(f)\ldots}_{cj(b)(r)\ldots} A^{(d)(r)}_{(l)(f)g} - \ldots, \end{cases}$$

- the M*-horizontal covariant derivative* of $D\Gamma(N)$:

$$(h_M) \quad \begin{cases} T^{ai(k)(d)\ldots}_{cj(b)(l)\ldots|s} = \dfrac{\delta T^{ai(k)(d)\ldots}_{cj(b)(l)\ldots}}{\delta x^s} + T^{fi(k)(d)\ldots}_{cj(b)(l)\ldots} H^a_{fs} + \\ + T^{ar(k)(d)\ldots}_{cj(b)(l)\ldots} H^i_{rs} + T^{ai(r)(d)\ldots}_{cj(f)(l)\ldots} H^{(f)(k)}_{(r)(b)s} + \ldots - \\ - T^{ai(k)(d)\ldots}_{fj(b)(l)\ldots} H^f_{cs} - T^{ai(k)(d)\ldots}_{cr(b)(l)\ldots} H^r_{js} - T^{ai(k)(f)\ldots}_{cj(b)(r)\ldots} H^{(d)(r)}_{(l)(f)s} - \ldots, \end{cases}$$

- the *vertical covariant derivative* of $D\Gamma(N)$:

$$(v)\quad \left\{\begin{array}{l} T^{ai(k)(d)\ldots}_{cj(b)(l)\ldots}|^{(s)}_{(g)} = \dfrac{\partial T^{ai(k)(d)\ldots}_{cj(b)(l)\ldots}}{\partial p^g_s} + T^{fi(k)(d)\ldots}_{cj(b)(l)\ldots} C^{a(s)}_{f(g)} + \\ + T^{ar(k)(d)\ldots}_{cj(b)(l)\ldots} C^{i(s)}_{r(g)} + T^{ai(r)(d)\ldots}_{cj(f)(l)\ldots} C^{(f)(k)(s)}_{(r)(b)(g)} + \ldots - \\ - T^{ai(k)(d)\ldots}_{fj(b)(l)\ldots} C^{f(s)}_{c(g)} - T^{ai(k)(d)\ldots}_{cr(b)(l)\ldots} C^{r(s)}_{j(g)} - T^{ai(k)(f)\ldots}_{cj(b)(r)\ldots} C^{(d)(r)(s)}_{(l)(f)(g)} - \ldots. \end{array}\right.$$

Remark 2.7 *If $T = Y$ is a d-vector field on E^*, locally expressed by*

$$Y = Y^a \frac{\delta}{\delta t^a} + Y^i \frac{\delta}{\delta x^i} + Y^{(a)}_{(i)} \frac{\partial}{\partial p^a_i},$$

then the following expressions of the local covariant derivatives hold good:

$$(h_{\mathcal{T}})\quad \left\{\begin{array}{l} Y^a_{\ /c} = \dfrac{\delta Y^a}{\delta t^c} + Y^b A^a_{bc}, \\ Y^i_{\ /c} = \dfrac{\delta Y^i}{\delta t^c} + Y^j A^i_{jc}, \\ Y^{(a)}_{(i)/c} = \dfrac{\delta Y^{(a)}_{(i)}}{\delta t^c} - Y^{(b)}_{(j)} A^{(a)(j)}_{(i)(b)c}, \end{array}\right. \qquad (h_M)\quad \left\{\begin{array}{l} Y^a_{\ |k} = \dfrac{\delta Y^a}{\delta x^k} + Y^b H^a_{bk}, \\ Y^i_{\ |k} = \dfrac{\delta Y^i}{\delta x^k} + Y^j H^i_{jk}, \\ Y^{(a)}_{(i)|k} = \dfrac{\delta Y^{(a)}_{(i)}}{\delta x^k} - Y^{(b)}_{(j)} H^{(a)(j)}_{(i)(b)k}, \end{array}\right.$$

$$(v)\quad \left\{\begin{array}{l} Y^a|^{(k)}_{(c)} = \dfrac{\partial Y^a}{\partial p^c_k} + Y^b C^{a(k)}_{b(c)}, \\ Y^i|^{(k)}_{(c)} = \dfrac{\partial Y^i}{\partial p^c_k} + Y^j C^{i(k)}_{j(c)}, \\ Y^{(a)}_{(i)}|^{(k)}_{(c)} = \dfrac{\partial Y^{(a)}_{(i)}}{\partial p^c_k} - Y^{(b)}_{(j)} C^{(a)(j)(k)}_{(i)(b)(c)}. \end{array}\right.$$

3 Components of h-normal N-linear connections on dual 1-jet spaces

Because the number of components which characterize an N-linear connection on E^* is big one (nine local components), we are constrained to study only a particular class of N-linear connections on E^*, which must be characterized by a reduced number of components. In this direction, let us fix on the temporal manifold $\mathcal{T}$ a semi-Riemannian metric h_{ab}, together with its Christoffel symbols χ^a_{bc}. Let $\mathbb{J}$ be the *h-normalization d-tensor field* on E^*, locally expressed by [5]

$$\mathbb{J} = J^{(i)}_{(a)bj} \delta p^a_i \otimes dt^b \otimes dx^j,$$

where $J^{(i)}_{(a)bj} = h_{ab}\delta^i_j$. In this context, we introduce the following geometrical concept:

Definition 3.1 *An N-linear connection $D\Gamma(N)$ on E^*, whose local components (2.3) verify the relations*

$$A^a_{bc} = \chi^a_{bc}, \qquad H^a_{bi} = 0, \qquad C^{a(i)}_{b(c)} = 0, \qquad D\mathbb{J} = 0,$$

*is called an h-**normal** N-**linear connection** on the dual 1-jet fibre bundle E^*.*

Theorem 3.2 *The adapted components of an h-normal N-linear connection $D\Gamma(N)$ verify the following identities:*

$$\begin{gathered} A^a_{bc} = \chi^a_{bc}, \quad H^a_{bi} = 0, \quad C^{a(i)}_{b(c)} = 0, \\ A^{(a)(j)}_{(i)(b)c} = \delta^a_b A^j_{ic} - \delta^j_i \chi^a_{bc}, \quad H^{(a)(j)}_{(i)(b)k} = \delta^a_b H^j_{ik}, \\ C^{(a)(j)(k)}_{(i)(b)(c)} = \delta^a_b C^{j(k)}_{i(c)}. \end{gathered} \tag{3.1}$$

Proof. It is obvious that the first three relations come immediately from the definition of an h-normal N-linear connection. To prove the other three relations, we emphasize that, taking into account the definition of the local $\mathcal{T}$-horizontal ("${}_{/g}$"), M-horizontal ("${}_{|s}$") and vertical ("$|^{(s)}_{(g)}$") covariant derivatives produced by $D\Gamma(N)$, the condition $D\mathbb{J} = 0$ is equivalent to

$$J^{(i)}_{(a)bj/g} = 0, \qquad J^{(i)}_{(a)bj|s} = 0, \qquad J^{(i)}_{(a)bj}|^{(s)}_{(g)} = 0.$$

Consequently, the condition $D\mathbb{J} = 0$ provides the local identities

$$\begin{gathered} h_{bf} A^{(f)(i)}_{(j)(a)c} = h_{ab} A^i_{jc} - \delta^i_j \left(\frac{\partial h_{ab}}{\partial t^c} - h_{ag} \chi^g_{bc} \right), \\ h_{bf} H^{(f)(i)}_{(j)(a)k} = h_{ba} H^i_{jk}, \quad h_{bf} C^{(f)(i)(k)}_{(j)(a)(c)} = h_{ba} C^{i(k)}_{j(c)}. \end{gathered}$$

Contracting now the above relations by h^{be}, we obtain the last required identities from (3.1). ■

Remark 3.3 *The above theorem says us that an h-normal N-linear connection on E^* is an N-linear connection determined by* ***four*** *effective components (instead of nine in the general case):*

$$D\Gamma(N) = \left(\chi^a_{bc}, \ A^i_{jc}, \ H^i_{jk}, \ C^{i(k)}_{j(c)} \right).$$

The other five components either vanish or are provided by the relations (3.1). Consequently, we can assert that the Berwald N_0-linear connection associated to the pair of metrics $\left(h_{ab}, \varphi_{ij}\right)$ is an h-normal N_0-linear connection on E^*, whose four effective components are

$$B\Gamma\left(N_0\right) = \left(\chi^a_{bc}, \ 0, \ \Gamma^i_{jk}, \ 0 \right).$$

4 Adapted components of torsion and curvature tensors

The study of the adapted components of the torsion and curvature tensors of an arbitrary N-linear connection $D\Gamma(N)$ on E^* was done in [5]. In that context, one proves that the torsion tensor $\mathbb{T}$ is determined by *twelve* effective local adapted d-tensors, while the curvature tensor $\mathbb{R}$ is determined by *eighteen* local adapted d-tensors. In what follows, we study the adapted components of the torsion and curvature tensors for an h-normal N-linear connection $D\Gamma(N)$.

Theorem 4.1 *The torsion tensor $\mathbb{T}$ of an h-normal N-linear connection $D\Gamma(N)$ is determined by **nine** effective local adapted d-tensors (instead of twelve in the general case):*

	h_T	h_M	v
h_Th_T	0	0	$R^{(f)}_{(r)ab}$
h_Mh_T	0	T^r_{aj}	$R^{(f)}_{(r)aj}$
vh_T	0	0	$P^{(f)\ (j)}_{(r)a(b)}$
h_Mh_M	0	T^r_{ij}	$R^{(f)}_{(r)ij}$
vh_M	0	$P^{r(j)}_{i(b)}$	$P^{(f)\ (j)}_{(r)i(b)}$
vv	0	0	$S^{(f)(i)(j)}_{(r)(a)(b)}$

(4.1)

where

$$T^r_{aj} = -A^r_{aj}, \quad T^r_{ij} = H^r_{ij} - H^r_{ji}, \quad P^{r(j)}_{i(b)} = C^{r(j)}_{i(b)},$$

$$P^{(f)\ (j)}_{(r)a(b)} = \frac{\partial \underset{1}{N}{}^{(f)}_{(r)a}}{\partial p^b_j} + \delta^f_b A^j_{ra} - \delta^j_r \chi^f_{ba}, \quad P^{(f)\ (j)}_{(r)i(b)} = \frac{\partial \underset{2}{N}{}^{(f)}_{(r)i}}{\partial p^b_j} + \delta^f_b H^j_{ri},$$

$$R^{(f)}_{(r)ab} = \frac{\delta \underset{1}{N}{}^{(f)}_{(r)a}}{\delta t^b} - \frac{\delta \underset{1}{N}{}^{(f)}_{(r)b}}{\delta t^a}, \quad R^{(f)}_{(r)aj} = \frac{\delta \underset{1}{N}{}^{(f)}_{(r)a}}{\delta x^j} - \frac{\delta \underset{2}{N}{}^{(f)}_{(r)i}}{\delta t^a},$$

$$R^{(f)}_{(r)ij} = \frac{\delta \underset{2}{N}{}^{(f)}_{(r)i}}{\delta x^j} - \frac{\delta \underset{2}{N}{}^{(f)}_{(r)j}}{\delta x^i}, \quad S^{(f)(i)(j)}_{(r)(a)(b)} = -\left(\delta^f_a C^{i(j)}_{r(b)} - \delta^f_b C^{j(i)}_{r(a)}\right).$$

Proof. Particularizing the general local expressions from [5], which generally give those twelve d-components of the torsion tensor of an N-linear connection, an h-normal N-linear connection $D\Gamma(N)$, we deduce that the adapted components T^a_{bc}, T^a_{bj} and $P^{a(k)}_{b(c)}$ vanish, while the other nine are given by the formulas from theorem. ■

Remark 4.2 *All torsion d-tensors of the Berwald h-normal N_0-linear connection $B\Gamma(N_0)$ (associated to the metrics h_{ab} and φ_{ij}) are zero, except*

$$R^{(f)}_{(r)ab} = \chi^f_{gab} p^g_r, \quad R^{(f)}_{(r)ij} = -\mathcal{R}^s_{rij} p^f_s,$$

where $\chi^f_{gab}(t)$ *(resp.* $\mathcal{R}^s_{rij}(x)$*) are the local curvature tensors of the semi-Riemannian metric* h_{ab} *(resp.* φ_{ij}*).*

Theorem 4.3 *The curvature tensor* $\mathbb{R}$ *of an* h*-normal* N*-linear connection* $D\Gamma(N)$ *is characterized by* ***seven*** *effective adapted local d-tensors (instead of eighteen in the general case):*

	$h_{\mathcal{T}}$	h_M	v
$h_{\mathcal{T}}h_{\mathcal{T}}$	χ^d_{abc}	R^l_{ibc}	$-R^{(d)(i)}_{(l)(a)bc} = \delta^i_l\chi^d_{abc} - \delta^d_a R^i_{lbc}$
$h_M h_{\mathcal{T}}$	0	R^l_{ibk}	$-R^{(d)(i)}_{(l)(a)bk} = -\delta^d_a R^i_{lbk}$
$wh_{\mathcal{T}}$	0	$P^{l\ (k)}_{ib(c)}$	$-P^{(d)(i)\ (k)}_{(l)(a)b(c)} = -\delta^d_a P^{i\ (k)}_{lb(c)}$
$h_M h_M$	0	R^l_{ijk}	$-R^{(d)(i)}_{(l)(a)jk} = -\delta^d_a R^i_{ljk}$
wh_M	0	$P^{l\ (k)}_{ij(c)}$	$-P^{(d)(i)\ (k)}_{(l)(a)j(c)} = -\delta^d_a P^{i\ (k)}_{lj(c)}$
ww	0	$S^{l(j)(k)}_{i(b)(c)}$	$-S^{(d)(i)(j)(k)}_{(l)(a)(b)(c)} = -\delta^d_a S^{i(j)(k)}_{l(b)(c)}$

(4.2)

where

$$R^d_{abc} := \chi^d_{abc} = \frac{\delta\chi^d_{ab}}{\delta t^c} - \frac{\delta\chi^d_{ac}}{\delta t^b} + \chi^f_{ab}\chi^d_{fc} - \chi^f_{ac}\chi^d_{fb},$$

$$R^l_{ibc} = \frac{\delta A^l_{ib}}{\delta t^c} - \frac{\delta A^l_{ic}}{\delta t^b} + A^r_{ib}A^l_{rc} - A^r_{ic}A^l_{rb} + C^{l(r)}_{i(f)}R^{(f)}_{(r)bc},$$

$$R^l_{ibk} = \frac{\delta A^l_{ib}}{\delta x^k} - \frac{\delta H^l_{ik}}{\delta t^b} + A^r_{ib}H^l_{rk} - H^r_{ik}A^l_{rb} + C^{l(r)}_{i(f)}R^{(f)}_{(r)bk},$$

$$P^{l\ (k)}_{ib(c)} = \frac{\partial A^l_{ib}}{\partial p^c_k} - C^{l(k)}_{i(c)/b} + C^{l(r)}_{i(f)}P^{(f)\ (k)}_{(r)b(c)},$$

$$R^l_{ijk} = \frac{\delta H^l_{ij}}{\delta x^k} - \frac{\delta H^l_{ik}}{\delta x^j} + H^r_{ij}H^l_{rk} - H^r_{ik}H^l_{rj} + C^{l(r)}_{i(f)}R^{(f)}_{(r)jk},$$

$$P^{l\ (k)}_{ij(c)} = \frac{\partial H^l_{ij}}{\partial p^c_k} - C^{l(k)}_{i(c)|j} + C^{l(f)}_{i(r)}P^{(r)\ (k)}_{(f)j(c)},$$

$$S^{l(j)(k)}_{i(b)(c)} = \frac{\partial C^{l(j)}_{i(b)}}{\partial p^c_k} - \frac{\partial C^{l(k)}_{i(c)}}{\partial p^b_j} + C^{r(j)}_{i(b)}C^{l(k)}_{r(c)} - C^{r(k)}_{i(c)}C^{l(j)}_{r(b)}.$$

Proof. The general formulas that express the local curvature d-tensors of an arbitrary N-linear connection (for more details, see [5]), applied to the particular case of an h-normal N-linear connection $D\Gamma(N)$, imply the above formulas and the relations from the Table (4.2). ■

Remark 4.4 *In the case of the Berwald h-normal* N_0*-linear connection* $B\Gamma(N_0)$ *(associated to the pair of metrics* $\big(h_{ab}, \varphi_{ij}\big)$*), all curvature d-tensors are zero, except*

$$R^d_{abc} = \chi^d_{abc}, \quad R^{(d)(i)}_{(l)(a)bc} = -\delta^i_l\chi^d_{abc}, \quad R^l_{ijk} = \mathcal{R}^l_{ijk} \quad R^{(d)(l)}_{(i)(a)jk} = \delta^d_a\mathcal{R}^l_{ijk},$$

where $\chi^f_{gab}(t)$ (resp. $\mathcal{R}^s_{rij}(x)$) are the local curvature tensors of the semi-Riemannian metric h_{ab} (resp. φ_{ij}).

5 Local Ricci identities. Non-metrical deflection d-tensor identities

Let us consider now the following more particular geometrical concept:

Definition 5.1 *An h-normal N-linear connection, whose local components*

$$C D\Gamma(N) = \left(\chi^a_{bc},\ A^i_{jc},\ H^i_{jk},\ C^{i(k)}_{j(c)}\right),$$

verify the relations

$$H^i_{jk} = H^i_{kj}, \qquad C^{i(k)}_{j(c)} = C^{k(i)}_{j(c)},$$

*is called an **h-normal N-linear connection of Cartan type** or a **$C D\Gamma(N)$-linear connection** on $E^* = J^{1*}(\mathcal{T}, M)$.*

Remark 5.2 *The torsion tensor $\mathbb{T}$ of an h-normal N-linear connection of Cartan type $C D\Gamma(N)$ is characterized only by **eight** adapted local d-tensors because the torsion components $T^i_{jk} = H^i_{jk} - H^i_{kj}$ from the Table (4.1) are vanishing.*

Example 5.3 *Taking into account that the Christoffel symbols $\Gamma^i_{jk}(x)$ of the spatial metric $\varphi_{ij}(x)$ are symmetric, it follows that the Berwald h-normal N_0-linear connection $B\Gamma(N_0)$ is of Cartan type.*

Theorem 5.4 *The following local **Ricci identities** for a $C D\Gamma(N)$-linear connection are true:*

- *the $h_{\mathcal{T}}$-Ricci identities:*

$$X^a_{/b/c} - X^a_{/c/b} = X^f \chi^a_{fbc} - X^a|^{(r)}_{(f)} R^{(f)}_{(r)bc},$$

$$X^a_{/b|k} - X^a_{|k/b} = -X^a_{|r} T^r_{bk} - X^a|^{(r)}_{(f)} R^{(f)}_{(r)bk},$$

$$X^a_{|j|k} - X^a_{|k|j} = -X^a|^{(r)}_{(f)} R^{(f)}_{(r)jk},$$

$$X^a_{/b}|^{(k)}_{(c)} - X^a|^{(k)}_{(c)/b} = -X^a|^{(r)}_{(f)} P^{(f)\ (k)}_{(r)b(c)},$$

$$X^a_{|j}|^{(k)}_{(c)} - X^a|^{(k)}_{(c)|j} = -X^a_{|r} C^{r(k)}_{j(c)} - X^a|^{(r)}_{(f)} P^{(f)\ (k)}_{(r)j(c)},$$

$$X^a|^{(j)}_{(b)}|^{(k)}_{(c)} - X^a|^{(k)}_{(c)}|^{(j)}_{(b)} = -X^a|^{(r)}_{(f)} S^{(f)(j)(k)}_{(r)(b)(c)};$$

- *the h_M-Ricci identities:*

$$X^i_{/b/c} - X^i_{/c/b} = X^r R^i_{rbc} - X^i|^{(r)}_{(f)} R^{(f)}_{(r)bc},$$

$$X^i_{/b|k} - X^i_{|k/b} = X^r R^i_{rbk} - X^i_{|r} T^r_{bk} - X^i|^{(r)}_{(f)} R^{(f)}_{(r)bk},$$

$$X^i_{|j|k} - X^i_{|k|j} = X^r R^i_{rjk} - X^i|^{(r)}_{(f)} R^{(f)}_{(r)jk},$$

$$X^i_{/b}|^{(k)}_{(c)} - X^i|^{(k)}_{(c)/b} = X^r P^{i\ \ (k)}_{rb(c)} - X^i|^{(r)}_{(f)} P^{(f)\ \ (k)}_{(r)b(c)},$$

$$X^i_{|j}|^{(k)}_{(c)} - X^i|^{(k)}_{(c)|j} = X^r P^{i\ \ (k)}_{rj(c)} - X^i_{|r} C^{r(k)}_{j(c)} - X^i|^{(r)}_{(f)} P^{(f)\ \ (k)}_{(r)j(c)},$$

$$X^i|^{(j)}_{(b)}|^{(k)}_{(c)} - X^i|^{(k)}_{(c)}|^{(j)}_{(b)} = X^r S^{i(j)(k)}_{r(b)(c)} - X^i|^{(r)}_{(f)} S^{(f)(j)(k)}_{(r)(b)(c)};$$

- *the v-Ricci identities:*

$$X^{(a)}_{(i)/b/c} - X^{(a)}_{(i)/c/b} = X^{(a)}_{(r)} R^r_{ibc} - X^{(f)}_{(i)} \chi^a_{fbc} - X^{(a)}_{(i)}|^{(r)}_{(f)} R^{(f)}_{(r)bc},$$

$$X^{(a)}_{(i)/b|k} - X^{(a)}_{(i)|k/b} = X^{(a)}_{(r)} R^r_{ibk} - X^{(a)}_{(i)|r} T^r_{bk} - X^{(a)}_{(i)}|^{(r)}_{(f)} R^{(f)}_{(r)bk},$$

$$X^{(a)}_{(i)|j|k} - X^{(a)}_{(i)|k|j} = X^{(a)}_{(r)} R^r_{ijk} - X^{(a)}_{(i)}|^{(r)}_{(f)} R^{(f)}_{(r)jk},$$

$$X^{(a)}_{(i)/b}|^{(k)}_{(c)} - X^{(a)}_{(i)}|^{(k)}_{(c)/b} = X^{(a)}_{(r)} P^{r\ \ (k)}_{ib(c)} - X^{(a)}_{(i)}|^{(r)}_{(f)} P^{(f)\ \ (k)}_{(r)b(c)},$$

$$X^{(a)}_{(i)|j}|^{(k)}_{(c)} - X^{(a)}_{(i)}|^{(k)}_{(c)|j} = X^{(a)}_{(r)} P^{r\ \ (k)}_{ij(c)} - X^{(a)}_{(i)|r} C^{r(k)}_{j(c)} - X^{(a)}_{(i)}|^{(r)}_{(f)} P^{(f)\ \ (k)}_{(r)j(c),}$$

$$X^{(a)}_{(i)}|^{(j)}_{(b)}|^{(k)}_{(c)} - X^{(a)}_{(i)}|^{(k)}_{(c)}|^{(j)}_{(b)} = X^{(a)}_{(r)} S^{r(j)(k)}_{i(b)(c)} - X^{(a)}_{(i)}|^{(r)}_{(f)} S^{(f)(j)(k)}_{(r)(b)(c)},$$

where

$$X = X^a \frac{\delta}{\delta t^a} + X^i \frac{\delta}{\delta x^i} + X^{(a)}_{(i)} \frac{\partial}{\partial p^a_i}$$

is an arbitrary d-vector field on the dual 1*-jet space* $E^* = J^{1*}(\mathcal{T}, M)$*.*

Proof. Let (Y_A) and (ω^A), where $A \in \left\{a, i, {}^{(a)}_{(i)}\right\}$, be on $E^* = J^{1*}(\mathcal{T}, M)$ the dual bases adapted to the nonlinear connection N, and let $X = X^F Y_F$ be a d-vector field on E^*. In this context, using the following true equalities (applied for a $CD\Gamma(N)$-linear connection D):

1. $D_{Y_C} Y_B = \Gamma^F_{BC} Y_F$,

2. $[Y_B, Y_C] = R^F_{BC} Y_F$,

3. $\mathbb{T}(Y_C, Y_B) = \mathbb{T}^F_{BC} Y_F = \{\Gamma^F_{BC} - \Gamma^F_{CB} - R^F_{CB}\} Y_F$,

4. $\mathbb{R}(Y_C, Y_B) Y_A = \mathbb{R}^F_{ABC} Y_F$,

5. $D_{Y_C} \omega^B = -\Gamma^B_{FC} \omega^F$,

6. $[\mathbb{R}(Y_C, Y_B) X] \otimes \omega^B \otimes \omega^C = \{D_{Y_C} D_{Y_B} X -$

   $- D_{Y_B} D_{Y_C} X - D_{[Y_C, Y_B]} X\} \otimes \omega^B \otimes \omega^C$,

by a direct calculation, we find that

$$X^A_{:B:C} - X^A_{:C:B} = X^F \mathbb{R}^A_{FBC} - X^A_{:F} \mathbb{T}^F_{BC}, \tag{5.1}$$

where "$_{:G}$" represents one from the local covariant derivatives "$_{/b}$", "$_{|j}$" or "$|^{(j)}_{(b)}$" produced by the h-normal N-linear connection of Cartan type $CD\Gamma(N)$.

Taking into account in (5.1) that the indices $A, B, C, \ldots$ belong to the set

$$\left\{a, i, {}^{(a)}_{(i)}\right\},$$

and using the particular features of an h-normal N-linear connection of Cartan type $CD\Gamma(N)$ (i.e., the torsion d-components T^i_{jk} are zero; we have the curvature relations from the Table (4.2)), by complicated computations, we find what we were looking for (see also the Table (4.1)). ∎

In order to find an interesting application of the preceding Ricci identities, let us consider the *canonical Liouville-Hamilton d-tensor field of polymomenta* on $E^* = J^{1*}(T, M)$, which is given by

$$\mathbb{C}^* = p^a_i \frac{\partial}{\partial p^a_i}.$$

In this context, for an h-normal N-linear connection of Cartan type $CD\Gamma(N)$, we can construct the *non-metrical deflection d-tensors*, setting

$$\Delta^{(a)}_{(i)b} = p^a_{i/b}, \quad \Delta^{(a)}_{(i)j} = p^a_{i|j}, \quad \vartheta^{(a)(j)}_{(i)(b)} = p^a_i|^{(j)}_{(b)},$$

where "$_{/b}$", "$_{|j}$" and "$|^{(j)}_{(b)}$" are the local covariant derivatives produced by $CD\Gamma(N)$.

By direct local computations, we deduce that the non-metrical deflection d-tensors of $CD\Gamma(N)$ have the expressions:

$$\Delta^{(a)}_{(i)b} = -\underset{1}{N}{}^{(a)}_{(i)b} - A^r_{ib} p^a_r + \chi^a_{fb} p^f_i, \quad \Delta^{(a)}_{(i)j} = -\underset{2}{N}{}^{(a)}_{(i)j} - H^r_{ij} p^a_r,$$
$$\vartheta^{(a)(j)}_{(i)(b)} = \delta^a_b \delta^j_i - C^{r(j)}_{i(b)} p^a_r.$$

Applying now the preceding (v)-set of Ricci identities (attached to an h-normal N-linear connection of Cartan type) to the components of the canonical Liouville-Hamilton d-vector field of polymomenta, we get

Corollary 5.5 *The following the **deflection d-tensor identities**, associated to an h-normal N-linear connection of Cartan type, are true:*

$$\left\{\begin{array}{l}
\Delta^{(a)}_{(i)b/c}-\Delta^{(a)}_{(i)c/b}=p^a_rR^r_{ibc}-p^f_i\chi^a_{fbc}-\vartheta^{(a)(r)}_{(i)(f)}R^{(f)}_{(r)bc}\\
\Delta^{(a)}_{(i)b|k}-\Delta^{(a)}_{(i)k/b}=p^a_rR^r_{ibk}-\Delta^{(a)}_{(i)r}T^r_{bk}-\vartheta^{(a)(r)}_{(i)(f)}R^{(f)}_{(r)bk}\\
\Delta^{(a)}_{(i)j|k}-\Delta^{(a)}_{(i)k|j}=p^a_rR^r_{ijk}-\vartheta^{(a)(r)}_{(i)(f)}R^{(f)}_{(r)jk}\\
\Delta^{(a)}_{(i)b}|^{(k)}_{(c)}-\vartheta^{(a)(k)}_{(i)(c)/b}=p^a_rP^{r\ \ (k)}_{ib(c)}-\vartheta^{(a)(r)}_{(i)(f)}P^{(f)\ \ (k)}_{(r)b(c)}\\
\Delta^{(a)}_{(i)j}|^{(k)}_{(c)}-\vartheta^{(a)(k)}_{(i)(c)|j}=p^a_rP^{r\ \ (k)}_{ij(c)}-\Delta^{(a)}_{(i)r}C^{r(k)}_{j(c)}-\vartheta^{(a)(r)}_{(i)(f)}P^{(f)\ (k)}_{(r)j(c)}\\
\vartheta^{(a)(j)}_{(i)(b)}|^{(k)}_{(c)}-\vartheta^{(a)(k)}_{(i)(c)}|^{(j)}_{(b)}=p^a_rS^{r(j)(k)}_{i(b)(c)}-\vartheta^{(a)(r)}_{(i)(f)}S^{(f)(j)(k)}_{(r)(b)(c)}.
\end{array}\right. \tag{5.2}$$

Remark 5.6 *The deflection d-tensor identities (5.2) will be used in the near future for the construction of the **geometrical Maxwell equations** that will govern the abstract multi-time geometrical "electromagnetism" produced by a quadratic Hamiltonian depending on polymomenta (this is our work in progress).*

6 The local Bianchi identities of the $CD\Gamma(N)$-connections on the dual jet bundle $J^{1*}(T,M)$

From the general theory of linear connections on a vector bundle, one knows that the torsions $\mathbb{T}$ and curvature $\mathbb{R}$ of a connection D on the dual 1-jet space $E^*=J^{1*}(T,M)$ are not independent. In other words, they are interrelated by the following general *Bianchi identities* (for any $X,Y,Z,U\in\mathcal{X}(E^*)$):

$$\sum_{\{X,Y,Z\}}\left\{(D_X\mathbb{T})(Y,Z)-\mathbb{R}(X,Y)Z+\mathbb{T}(\mathbb{T}(X,Y),Z)\right\}=0,$$
$$\sum_{\{X,Y,Z\}}(D_X\mathbb{R})(Y,Z,U)+\mathbb{R}(\mathbb{T}(X,Y),Z)U=0,$$

where $\Sigma_{\{X,Y,Z\}}$ means a cyclic sum. Obviously, working with a $CD\Gamma(N)$-linear connection and the local adapted basis of d-vector fields $(X_A)\subset\mathcal{X}(E^*)$ (associated to the given nonlinear connection N on E^*), the above Bianchi identities are locally described by the equalities:

$$\begin{array}{c}
\sum\limits_{\{A,B,C\}}\left\{\mathbb{R}^F_{ABC}-\mathbb{T}^F_{AB:C}-\mathbb{T}^G_{AB}\mathbb{T}^F_{CG}\right\}=0,\\
\sum\limits_{\{A,B,C\}}\left\{\mathbb{R}^F_{DAB:C}+\mathbb{T}^G_{AB}\mathbb{R}^F_{DCG}\right\}=0,
\end{array} \tag{6.1}$$

where $\mathbb{R}(X_A,X_B)X_C=\mathbb{R}^D_{CBA}X_D$, $\mathbb{T}(X_A,X_B)=\mathbb{T}^D_{BA}X_D$, and "$_{:C}$" represents one from the local covariant derivatives "$_{/a}$", "$_{|i}$" or "$|^{(i)}_{(a)}$" of the $CD\Gamma(N)$-linear connection D (for similar details, see the works [14], [15] and [17]). Consequently, we find:

Theorem 6.1 *The following **thirty** effective local **Bianchi identities** for an h-normal N-linear connection of Cartan type $C\Gamma(N)$ are true on the dual 1-jet space $E^* = J^{1*}(\mathcal{T}, M)$:*

- *the first set:*

  1. $\sum_{\{a,b,c\}} \chi^d_{abc} = 0,$
  2. $\mathcal{A}_{\{a,b\}} \left\{ T^l_{ar} T^r_{bk} - T^l_{ak/b} \right\} = R^l_{kab} - C^{l(r)}_{k(f)} R^{(f)}_{(r)ab},$
  3. $\mathcal{A}_{\{j,k\}} \left\{ C^{l(r)}_{k(f)} R^{(f)}_{(r)aj} + R^l_{jak} + T^l_{aj|k} \right\} = 0,$
  4. $\sum_{\{i,j,k\}} \left\{ C^{l(r)}_{k(f)} R^{(f)}_{(r)ij} - R^l_{ijk} \right\} = 0,$

- *the second set:*

  5. $\sum_{\{a,b,c\}} \left\{ R^{(d)}_{(l)ab/c} + P^{(d)\ \ (r)}_{(l)c(f)} R^{(f)}_{(r)ab} \right\} = 0,$
  6. $\mathcal{A}_{\{a,b\}} \left\{ R^{(d)}_{(l)ak/b} + P^{(d)\ \ (r)}_{(l)b(f)} R^{(f)}_{(r)ak} + R^{(d)}_{(l)br} T^r_{ak} \right\} =$

     $= R^{(d)}_{(l)ab|k} + P^{(d)\ \ (r)}_{(l)k(f)} R^{(f)}_{(r)ab},$
  7. $\mathcal{A}_{\{j,k\}} \left\{ R^{(d)}_{(l)aj|k} + P^{(d)\ \ (r)}_{(l)k(f)} R^{(f)}_{(r)aj} + R^{(d)}_{(l)kr} T^r_{aj} \right\} =$

     $= -R^{(d)}_{(l)jk/a} - P^{(d)\ \ (r)}_{(l)a(f)} R^{(f)}_{(r)jk},$
  8. $\sum_{\{i,j,k\}} \left\{ R^{(d)}_{(l)ij|k} + P^{(d)\ \ (r)}_{(l)k(f)} R^{(f)}_{(r)ij} \right\} = 0,$

- *the third set:*

  9. $T^l_{ak}|^{(p)}_{(e)} - C^{l(p)}_{r(e)} T^r_{ak} + P^{l\ \ (p)}_{ka(e)} + C^{l(p)}_{k(e)/a} - C^{l(r)}_{k(f)} P^{(f)\ \ (p)}_{(r)a(e)} + C^{r(p)}_{k(e)} T^l_{ar} = 0,$
  10. $\mathcal{A}_{\{j,k\}} \left\{ C^{l(p)}_{j(e)|k} + C^{l(r)}_{k(f)} P^{(f)\ \ (p)}_{(r)j(e)} + P^{l\ \ (p)}_{jk(e)} \right\} = 0,$

- *the fourth set:*

  11. $\mathcal{A}_{\{a,b\}} \left\{ P^{(d)\ \ (p)}_{(l)a(e)/b} + P^{(d)\ \ (r)}_{(l)b(f)} P^{(f)\ \ (p)}_{(r)a(e)} \right\} =$

      $= R^{(d)}_{(l)ab}|^{(p)}_{(e)} + R^{(d)(p)}_{(l)(e)ab} + S^{(d)(p)(r)}_{(l)(e)(f)} R^{(f)}_{(r)ab},$

12. $\mathcal{A}_{\{a,k\}}\left\{P^{(d)\;\;(p)}_{(l)a(e)|k}+P^{(d)\;\;(r)}_{(l)k(f)}P^{(f)\;\;(p)}_{(r)a(e)}\right\}=$

$$=R^{(d)}_{(l)ak}\Big|^{(p)}_{(e)}+R^{(d)(p)}_{(l)(e)ak}+S^{(d)(p)(r)}_{(l)(e)(f)}R^{(f)}_{(r)ak}+R^{(d)}_{(l)ar}C^{r(p)}_{k(e)}-T^{r}_{ak}P^{(d)\;(p)}_{(l)r(e)}$$

13. $\mathcal{A}_{\{j,k\}}\left\{P^{(d)\;(p)}_{(l)j(e)|k}+P^{(d)\;(r)}_{(l)k(f)}P^{(f)\;(p)}_{(r)j(e)}+R^{(d)}_{(l)kr}C^{r(p)}_{j(e)}\right\}=$

$$=R^{(d)}_{(l)jk}\Big|^{(p)}_{(e)}+R^{(d)(p)}_{(l)(e)jk}+S^{(d)(p)(r)}_{(l)(e)(f)}R^{(f)}_{(r)jk},$$

- *the fifth set:*

14. $\mathcal{A}_{\left\{{}^{(j)}_{(b)},{}^{(k)}_{(c)}\right\}}\left\{C^{l(j)}_{i(b)}\Big|^{(k)}_{(c)}+C^{r(k)}_{i(c)}C^{l(j)}_{r(b)}\right\}=S^{l(j)(k)}_{i(b)(c)}-C^{l(r)}_{i(f)}S^{(f)(j)(k)}_{(r)(b)(c)},$

- *the sixth set:*

15. $\mathcal{A}_{\left\{{}^{(j)}_{(b)},{}^{(k)}_{(c)}\right\}}\left\{P^{(d)\;\;(j)}_{(l)a(b)}\Big|^{(k)}_{(c)}+P^{(f)\;\;(j)}_{(r)a(b)}S^{(d)(k)(r)}_{(l)(c)(f)}-P^{(d)(j)\;\;(k)}_{(l)(b)a(c)}\right\}=$

$$=-S^{(d)(j)(k)}_{(l)(b)(c)/a}-S^{(f)(j)(k)}_{(r)(b)(c)}P^{(d)\;\;(r)}_{(l)a(f)},$$

16. $\mathcal{A}_{\left\{{}^{(j)}_{(b)},{}^{(k)}_{(c)}\right\}}\left\{P^{(d)\;(j)}_{(l)i(b)}\Big|^{(k)}_{(c)}+P^{(f)\;(j)}_{(r)i(b)}S^{(d)(k)(r)}_{(l)(c)(f)}-P^{(d)(j)\;(k)}_{(l)(b)i(c)}-\right.$

$$\left.-C^{r(j)}_{i(b)}P^{(d)\;\;(k)}_{(l)r(c)}\right\}=-S^{(d)(j)(k)}_{(l)(b)(c)|i}-S^{(f)(j)(k)}_{(r)(b)(c)}P^{(d)\;(r)}_{(l)i(f)},$$

- *the seventh set:*

17. $\sum_{\left\{{}^{(i)}_{(a)},{}^{(j)}_{(b)},{}^{(k)}_{(c)}\right\}}\left\{S^{(d)(i)(j)}_{(l)(a)(b)}\Big|^{(k)}_{(c)}+S^{(f)(i)(j)}_{(r)(a)(b)}S^{(d)(k)(r)}_{(l)(c)(f)}+S^{(d)(i)(j)(k)}_{(l)(a)(b)(c)}\right\}=0,$

- *the eight set:*

18. $\sum_{\{a,b,c\}}\chi^{d}_{eab/c}=0,$

19. $\chi^{d}_{eab|k}=0,$

20. $\chi^{d}_{eab}\Big|^{(k)}_{(c)}=0,$

21. $\sum_{\{a,b,c\}}\left\{R^{l}_{pab/c}+R^{(f)}_{(r)ab}P^{l\;\;(r)}_{pc(f)}\right\}=0,$

22. $\mathcal{A}_{\{a,b\}}\left\{R^{l}_{pak/b}+R^{(f)}_{(r)ak}P^{l\;\;(r)}_{pb(f)}+T^{r}_{ak}R^{l}_{pbr}\right\}=R^{l}_{pab|k}+R^{(f)}_{(r)ab}P^{l\;\;(r)}_{pk(f)},$

23. $\mathcal{A}_{\{j,k\}}\left\{R^{l}_{paj|k}+R^{(f)}_{(r)aj}P^{l\ \ (r)}_{pk(f)}+T^{r}_{aj}R^{l}_{pkr}\right\}=-R^{l}_{pjk/a}-R^{(f)}_{(r)jk}P^{l\ \ (r)}_{pa(f)}$,

24. $\sum_{\{i,j,k\}}\left\{R^{l}_{pij|k}+R^{(f)}_{(r)ij}P^{l\ \ (r)}_{pk(f)}\right\}=0$,

- *the nineth set:*

25. $\mathcal{A}_{\{a,b\}}\left\{P^{l\ \ (p)}_{ia(e)/b}+P^{(f)\ \ (p)}_{(r)a(e)}P^{l\ (r)}_{ib(f)}\right\}=R^{l}_{iab}|^{(p)}_{(e)}+R^{(f)}_{(r)ab}S^{l(p)(r)}_{i(e)(f)}$,

26. $\mathcal{A}_{\{a,k\}}\left\{P^{l\ \ (p)}_{ia(e)|k}+P^{(f)\ \ (p)}_{(r)a(e)}P^{l\ \ (r)}_{ik(f)}\right\}=$

    $=R^{l}_{iak}|^{(p)}_{(e)}+R^{(f)}_{(r)ak}S^{l(p)(r)}_{i(e)(f)}+C^{r(p)}_{k(e)}R^{l}_{iar}-T^{r}_{ak}P^{l\ \ (p)}_{ir(e)}$,

27. $\mathcal{A}_{\{j,k\}}\left\{P^{l\ \ (p)}_{ij(e)|k}+P^{(f)\ \ (p)}_{(r)j(e)}P^{l\ \ (r)}_{ik(f)}+C^{r(p)}_{j(e)}R^{l}_{ikr}\right\}=$

    $=R^{l}_{ijk}|^{(p)}_{(e)}+R^{(f)}_{(r)jk}S^{l(p)(r)}_{i(e)(f)}$,

- *the tenth set:*

28. $\mathcal{A}_{\left\{\binom{(j)}{(b)},\binom{(k)}{(c)}\right\}}\left\{P^{l\ \ (j)}_{pa(b)}|^{(k)}_{(c)}+P^{(f)\ \ (j)}_{(r)a(b)}S^{l(k)(r)}_{p(c)(f)}\right\}=$

    $=-S^{l(j)(k)}_{p(b)(c)/a}-S^{(f)(j)(k)}_{(r)(b)(c)}P^{l\ \ (r)}_{pa(f)}$,

29. $\mathcal{A}_{\left\{\binom{(j)}{(b)},\binom{(k)}{(c)}\right\}}\left\{P^{l\ \ (j)}_{pi(b)}|^{(k)}_{(c)}+P^{(f)\ \ (j)}_{(r)i(b)}S^{l(k)(r)}_{p(c)(f)}-C^{r(j)}_{i(b)}P^{l\ \ (k)}_{pr(c)}\right\}=$

    $=-S^{l(j)(k)}_{p(b)(c)|i}-S^{(f)(j)(k)}_{(r)(b)(c)}P^{l\ \ (r)}_{pi(f)}$,

- *the eleventh set:*

30. $\sum_{\left\{\binom{(i)}{(a)},\binom{(j)}{(b)},\binom{(k)}{(c)}\right\}}\left\{S^{l(i)(j)}_{p(a)(b)}|^{(k)}_{(c)}+S^{(r)(i)(j)}_{(f)(a)(b)}S^{l(k)(f)}_{p(c)(r)}\right\}=0$,

where, if $\{A,B,C\}$ *are indices of type* $\left\{a,i,\binom{(a)}{(i)}\right\}$, *then* $\sum_{\{A,B,C\}}$ *represents a cyclic sum, and* $\mathcal{A}_{\{A,B\}}$ *represents an alternate sum.*

Proof. Taking into account that the indices $A,B,C,D...$ are of type

$$\left\{a,i,\binom{(a)}{(i)}\right\},$$

and the torsion $\mathbb{T}^{C}_{AB}$ and curvature $\mathbb{R}^{D}_{ABC}$ adapted components are given in the Tables (4.1) and (4.2), after laborious local computations, the formulas (6.1) imply the required Bianchi identities. ■

Remark 6.2 *We point out that, in the particular single-time case*

$$(\mathcal{T}, h) = (\mathbb{R}, \delta = 1),$$

the last identity of our each set of local Bianchi identities reduces to one of the classical eleven Bianchi identities that characterize the N*-linear connections in the classical Hamilton geometry on cotangent bundles (see* [15]*).*

Acknowledgements. The authors of this paper would like to express their sincere gratitude to Professor Gh. Atanasiu for his suggestions and useful discussions on this research topic.

References

[1] G.S. Asanov, *Jet extension of Finslerian gauge approach*, Fortschritte der Physik, vol. **38**, no. **8** (1990), 571-610.

[2] Gh. Atanasiu, *The invariant expression of Hamilton geometry*, Tensor N.S., vol. **47**, no. **3** (1988), 225-234.

[3] Gh. Atanasiu, F.C. Klepp, *Nonlinear connections in cotangent bundle*, Publ. Math. Debrecen, Hungary, vol. **39**, no. **1-2** (1991), 107-111.

[4] Gh. Atanasiu, M. Neagu, *Distinguished tensors and Poisson brackets in the multi-time Hamilton geometry*, BSG Proceedings **16**, Geometry Balkan Press, Bucharest (2009), 12-27.

[5] Gh. Atanasiu, M. Neagu, *Distinguished torsion, curvature and deflection tensors in the multi-time Hamilton geometry*, Electronic Journal "Differential Geometry-Dynamical Systems", vol. **11** (2009), 20-40.

[6] G. Giachetta, L. Mangiarotti, G. Sardanashvily, *Covariant Hamiltonian field theory*, http://arXiv.org/hep-th/9904062 (1999).

[7] G. Giachetta, L. Mangiarotti, G. Sardanashvily, *Polysymplectic Hamiltonian formalism and some quantum outcomes*, http://arXiv.org/hep-th/0411005 (2004).

[8] M. Gotay, J. Isenberg, J.E. Marsden, R. Montgomery, *Momentum maps and classical fields. Part I. Covariant field theory*, http://arXiv.org/physics/9801019 (2004).

[9] M. Gotay, J. Isenberg, J.E. Marsden, *Momentum maps and classical fields. Part II. Canonical analysis of field theories*, http://arXiv.org/math-ph/0411032 (2004).

[10] I.V. Kanatchikov, *Basic structures of the covariant canonical formalism for fields based on the De Donder-Weyl theory*, http://arXiv.org/hep-th/9410238 (1994).

[11] I.V. Kanatchikov, *On quantization of field theories in polymomentum variables*, AIP Conf. Proc., vol. **453**, Issue **1** (1998), 356-367.

[12] I.V. Kanatchikov, *On the canonical structure of the De Donder-Weyl covariant Hamiltonian formulation of field theory I. Graded Poisson brackets and equations of motion*, http://arXiv.org/hep-th/9312162 (1993).

[13] R. Miron, *Hamilton geometry*, An. Şt. "Al. I. Cuza" Univ., Iaşi, Romania, vol. **35** (1989), 33-67.

[14] R. Miron, M. Anastasiei, *The Geometry of Lagrange Spaces: Theory and Applications*, Kluwer Academic Publishers, 1994.

[15] R. Miron, D. Hrimiuc, H. Shimada, S.V. Sabău, *The Geometry of Hamilton and Lagrange Spaces*, Kluwer Academic Publishers, 2001.

[16] R. Miron, M.S. Kirkovits, M. Anastasiei, *A geometrical model for variational problems of multiple integrals*, Proc. Conf. Diff. Geom. and Appl., Dubrovnik, Yugoslavia, June 26-July 3, (1988), 8-25.

[17] M. Neagu, *Ricci and Bianchi identities for h-normal Γ-linear connections on $J^1(T, M)$*, Hindawi Publishing Corporation, International Journal of Mathematics and Mathematical Sciences, no. **34** (2003), 2177-2192.

[18] M. Neagu, *Riemann-Lagrange Geometry on* 1*-Jet Spaces*, Matrix Rom, Bucharest, 2005.

[19] M. Neagu, C. Udrişte, *Torsion, curvature and deflection d-tensors on* $J^1(T, M)$, Balkan Journal of Geometry and Its Applications, vol. **6**, no. **1** (2001), 29-45.

[20] M. Neagu, C. Udrişte, A. Oană, *Multi-time dependent sprays and h-traceless maps,* Balkan Journal of Geometry and Its Applications, vol. **10**, no. **2** (2005), 76-92.

[21] P.J. Olver, *Applications of Lie Groups to Differential Equations*, Graduate Texts in Mathematics, vol. **107**, Springer-Verlag, New York, 1986.

[22] D.J. Saunders, *The Geometry of Jet Bundles*, Cambridge University Press, New York, London, 1989.

Alexandru OANĂ and Mircea NEAGU
University Transilvania of Braşov,
Department of Mathematics - Informatics,
Blvd. Iuliu Maniu, no. 50, Braşov 500091, Romania.
E-mails: alexandru.oana@unitbv.ro, mircea.neagu@unitbv.ro